

DISORDERED BOSE EINSTEIN CONDENSATES WITH INTERACTION

ROBERT SEIRINGER, JAKOB YNGVASON, AND VALENTIN A. ZAGREBNOV

ABSTRACT. We study the effects of random scatterers on the ground state of the one-dimensional Lieb-Liniger model of interacting bosons on the unit interval in the Gross-Pitaevskii regime. We prove that Bose Einstein condensation survives even a strong random potential with a high density of scatterers. The character of the wave function of the condensate, however, depends in an essential way on the interplay between randomness and the strength of the two-body interaction. For low density of scatterers or strong interactions the wave function extends over the whole interval. High density of scatterers and weak interaction, on the other hand, leads to localization of the wave function in a fragmented subset of the interval.

1. INTRODUCTION

While the effects of random potentials on single particle Schrödinger operators [1] and ideal Bose gases^{2–4} are rather well explored, the present understanding of such effects on many-body systems of interacting particles is much less complete. In recent years, however, many papers concerning the interplay of Bose-Einstein Condensation (BEC) and disorder have appeared of which references [5]–[16] are but a sample.

We present here results on a model that is in a sense the simplest one imaginable where this interplay can be studied by rigorous mathematical means. This is the one-dimensional Lieb-Liniger (LL) model [17] of bosons with contact interaction in a ‘flat’ trap, augmented by an external random potential that is generated by Poisson distributed point scatterers of equal strength. We study the ground state and prove that, no matter the strength of the random potential, BEC is not destroyed by the random potential in the Gross-Pitaevskii (GP) limit where the particle number tends to infinity while the coupling parameter in a mean-field scaling stays fixed. The character of the wave function of the condensate, however, depends in an essential way on the relative size of the three parameters involved. These are the scaled coupling parameter γ for the interaction among the particles, the density of the scatterers ν , and the strength σ of the scattering potential. All these parameters are assumed to be large in suitable units.

Our main results can be summarized as follows: For $\sigma \gg 1$ we identify three different “phases” of the condensate. For large $\gamma \gg \nu^2$ the condensate is extended over the whole trap (the unit interval in our model). A transition from a delocalized

Key words and phrases. Bose Einstein condensation; random Hamiltonians; one-dimensional models.

Contribution to the proceedings of ICMP12, Aalborg, Denmark, August 6–11, 2012.

© 2012 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

to a localized state takes place when γ is of the order ν^2 , in the sense that for $\gamma \ll \nu^2$ the density is essentially distributed among a fraction $\lambda \ll 1$ of the $\nu \gg 1$ intervals between the obstacles. For $\gamma \gg \nu/(\ln \nu)^2$ we still have $\lambda \nu \gg 1$, but for $\gamma \sim \nu/(\ln \nu)^2$ the fraction of intervals that are significant occupied shrinks to $O(\nu^{-1})$. We stress, however, that in all cases there is complete BEC into a single state in the limit when the particle number tends to infinity.

Details of our analysis are given in the preprint arXiv:1207.7054.

2. THE MODEL

The model we consider is the Lieb-Liniger model of bosons with contact interaction on the unit interval but with an additional external random potential V_ω . The Hamiltonian on the Hilbert space $L^2([0, 1], dz)^{\otimes_s N}$ is

$$(1) \quad H = \sum_{i=1}^N (-\partial_{z_i}^2 + V_\omega(z_i)) + \frac{\gamma}{N} \sum_{i < j} \delta(z_i - z_j)$$

with $\gamma \geq 0$ and Dirichlet boundary conditions. The random potential is taken to be

$$(2) \quad V_\omega(z) = \sigma \sum_j \delta(z - z_j^\omega)$$

with $\sigma \geq 0$ independent of the random sample ω while the *obstacles* $\{z_j^\omega\}$ are *Poisson distributed* with density $\nu \gg 1$, i.e., their mean distance is ν^{-1} .

There are several reasons for studying this model:

- It is the simplest model of its kind.
- BEC in the ground state can be proved in a suitable limit, for an *arbitrary* nonnegative external potential.
- V_ω is simple enough to allow a rigorous analysis of the condensate.

We remark also that the case $\gamma = 0$, $\sigma = \infty$ corresponds to the *Luttinger-Sy model* $\square$.

Once BEC has been established the main question concerns the dependence of the properties of the condensate on the three parameters γ , σ and ν .

3. BEC IN THE GROUND STATE

A basic fact about the model (1) is that for *fixed* γ , σ and configuration ω there is *complete BEC in the ground state* in the sense that the 1-particle density matrix/ N converges to a one dimensional projector as $N \rightarrow \infty$. The corresponding wave function of the condensate is the L^2 -normalized minimizer of the *Gross-Pitaevskii (GP) energy functional*

$$(3) \quad \mathcal{E}^{\text{GP}}[\psi] = \int_0^1 \{ |\psi'(z)|^2 + V_\omega(z) |\psi(z)|^2 + (\gamma/2) |\psi(z)|^4 \} dz.$$

We want, however, to consider ν , σ and γ large. Hence it is important to estimate also the rate of the convergence of the 1-particle density matrix as $N \rightarrow \infty$, in dependence of these parameters and of the configurations ω .

3.1. **The Proof of BEC (sketch).** The proof of BEC is based on energy bounds:

- An upper bound to the many-body ground state energy E_0^{QM} by taking $\psi_0^{\otimes N}$ as a trial function for H where ψ_0 is the minimizer of the GP energy functional, normalized so that $\|\psi_0\|_2 = 1$. This gives

$$(4) \quad E_0^{\text{QM}} \leq N e_0$$

where $e_0 = \mathcal{E}^{\text{GP}}[\psi_0]$ is the g.s.e. of the GP functional.

- An *operator lower bound* for the many-body Hamiltonian H , up to controlled errors, in terms of the 1-particle mean-field Hamiltonian

$$(5) \quad h = -\partial_z^2 + V_\omega(z) + \gamma|\psi_0(z)|^2 - (\gamma/2)\int|\psi_0|^4$$

which has ψ_0 as ground state with energy e_0 .

BEC follows from the upper and lower bounds and the fact that there is an energy gap between e_0 and the next lowest eigenvalue, e_1 , of the mean-field Hamiltonian h :

Let

$$(6) \quad N_0 = \langle \Psi_0 | a^\dagger(\psi_0) a(\psi_0) | \Psi_0 \rangle$$

be the occupation number of the GP ground state ψ_0 in the many-body ground state Ψ_0 . Then the energy bounds give

$$(7) \quad N_0 e_0 + (N - N_0) e_1 - o(1) N e_0 \leq E_0^{\text{QM}} \leq N e_0$$

where the $o(1)$ factor depends only on γ and N . This implies an upper bound for the depletion:

$$(8) \quad \left(1 - \frac{N_0}{N}\right) \leq o(1) \frac{e_0}{e_1 - e_0}$$

More precisely, the estimate on the depletion of the condensate is

THEOREM 1 (BEC).

$$(9) \quad \left(1 - \frac{N_0}{N}\right) \leq C \frac{e_0}{e_1 - e_0} N^{-1/3} \min\{\gamma^{1/2}, \gamma\}.$$

We must, however, also consider the dependence of the energy gap, $e_1 - e_0$, on the random potential and the parameters.

3.2. **Remark.** If $N_{\leq k}$ denotes the occupation of the k lowest eigenvalues of (5) with energies up to e_k , then we have

$$(10) \quad \left(1 - \frac{N_{\leq k}}{N}\right) \leq C \frac{e_0}{e_k - e_0} N^{-1/3} \min\{\gamma^{1/2}, \gamma\}.$$

For finite N this estimate may be more useful than (9) because even if $k \ll N$, $e_k - e_0$ can be substantially larger than $e_1 - e_0$. Thus the right side of (10) can be < 1 even in cases when (9) contains no information because the right-hand side is > 1 .

3.3. The energy gap. Consider a one-dimensional Schrödinger operator $-\partial_z^2 + W(z)$ on the unit interval with a nonnegative potential W and Dirichlet boundary conditions.

LEMMA 1 (Gap). *Define $\eta > 0$ by $\eta^2 = \pi^2 + 3 \int_0^1 W(z) dz$. Then*

$$(11) \quad e_1 - e_0 \geq \eta \ln(1 + \pi e^{-2\eta})$$

The proof is based on a modification of a result of Kirsch and Simon [19], that involves the sup norm of W instead of the integral. In our case $\eta = \eta_\omega = \sqrt{\pi^2 + 3m_\omega\sigma + 3\gamma}$ where m_ω is the number of obstacles in $[0, 1]$, that is almost surely equal to ν in the limit $\nu \rightarrow \infty$.

For large σm_ω the bound is certainly not optimal, in fact, in this case one expects $e_1 - e_0 \sim (\sigma m_\omega)^{-1}$.

3.4. The Poisson distribution of the obstacles. Let $z_1^\omega \leq z_2^\omega \leq \dots \leq z_{m_\omega}^\omega$ denote those random points which lie in the unit interval $[0, 1]$. The lengths $\ell_i = z_{i+1}^\omega - z_i^\omega$ are independent random variables with distribution

$$(12) \quad dP_\nu(\ell) = \nu e^{-\ell\nu} d\ell$$

and we are considering the case $\nu \gg 1$.

The average length of an interval free of obstacles is ν^{-1} and with probability one, $m_\omega/\nu \rightarrow 1$ and $\sum_i \ell_i = 1$ for $\nu \gg 1$. Combined with the estimate on the energy gap this implies that the depletion of the BEC is uniform in the L^p norm on sample space for any $p < \infty$.

3.5. Ideal vs interacting gas. While the average length of an interval is ν^{-1} there is, with probability one, a unique largest interval of length $\sim \nu^{-1} \ln \nu$. If there is no interaction, i.e., $\gamma = 0$, and $\sigma \rightarrow \infty$ (Luttinger-Sy model), then the ground state wave function will be localized in the largest interval for kinetic energy reasons, with energy $\sim \nu^2/(\ln \nu)^2$.

The question is how the character of the wave function of the condensate, ψ_0 , changes when the interaction, i.e., the term $(\gamma/N) \int |\psi_0(z)|^4 dz$ comes into play.

4. THE MASS DISTRIBUTION OF THE CONDENSATE

4.1. A limit theorem for the GP energy. Our first result is that the energy becomes *deterministic* in an appropriate limit.

Let $e_\omega(\gamma, \sigma, \nu)$ denote the GP energy with the random potential V_ω and $e_0(\gamma, \nu)$ the energy for $\sigma = \infty$, averaged over ω .

THEOREM 2 (Convergence of the energy). *Assume that $\nu \rightarrow \infty$, $\sigma \rightarrow \infty$ and $\gamma \rightarrow \infty$ in such a way that*

$$(13) \quad \gamma \gg \frac{\nu}{(\ln \nu)^2} \quad \text{and} \quad \sigma \gg \frac{\nu}{1 + \ln(1 + \nu^2/\gamma)}.$$

Then, for almost every sample ω ,

$$(14) \quad \lim \frac{e_\omega(\gamma, \sigma, \nu)}{e_0(\gamma, \nu)} = 1.$$

4.2. Comments on the proof. The proof has two interrelated parts:

- Comparison of $e_\omega(\gamma, \sigma, \nu)$ with $e_\omega(\gamma, \infty, \nu)$.
- Comparison of $e_\omega(\gamma, \infty, \nu)$ with the deterministic $e_0(\gamma, \nu)$.

Calculations are conveniently done in a grand canonical ensemble, introducing a chemical potential μ that determines the optimal repartition

$$(15) \quad n(\ell) \approx (\ell\gamma)^{-1} [\mu\ell^2 - \pi^2]_+$$

of the condensate mass in intervals of length ℓ between the obstacles. The lengths are distributed according to $dP_\nu(\ell)$ and the normalization requires

$$(16) \quad \nu \int n(\ell) dP_\nu(\ell) = 1.$$

4.3. Discussion of the GP wave function for large σ . For large σ the average number of intervals with non-zero occupation numbers is given by

$$(17) \quad \nu \int_{\pi/\sqrt{\mu}}^{\infty} dP_\nu(\ell) = e^{-\pi\nu/\sqrt{\mu}} \nu.$$

Since ν is the total (average) number of available intervals,

$$(18) \quad \lambda := e^{-\pi\nu/\sqrt{\mu}} \leq 1$$

defines the fraction of them which are occupied. The normalization requires

$$(19) \quad 1 \sim \frac{\mu}{\gamma} e^{-\pi\nu/\sqrt{\mu}}$$

so λ is determined by γ and ν via relation

$$(20) \quad \frac{\lambda}{(\ln \lambda^{-1})^2} \sim \frac{\gamma}{\nu^2}.$$

We can now distinguish the following *limiting cases*:

- If $\gamma \gg \nu^2$ then by (20) we get $\lambda \rightarrow 1$, i.e., *all* the intervals are occupied (delocalization). The chemical potential satisfies $\mu \sim \gamma$ in this regime.
- If $\gamma \sim \nu^2$ then $\lambda \sim 1$, but λ is strictly less than 1 (transition to localization). Again we have $\mu \sim \gamma$.
- If $\gamma \ll \nu^2$ then $\lambda \ll 1$, i.e., only a small fraction of the intervals are occupied (localization). The relation (19) implies $\mu \sim \frac{\gamma}{\lambda}$ for the chemical potential.
- If $\gamma \sim \nu/(\ln \nu)^2$ then by (20) the fraction λ becomes $O(1/\nu)$, i.e., only finitely many intervals are occupied. In this latter case, $\mu \sim \gamma\nu \sim \nu^2/(\ln \nu)^2$, which corresponds exactly to the inverse of the square of the size of the *largest* interval.

In particular, $\lambda\nu \gg 1$ only if $\gamma \gg \nu/(\ln \nu)^2$, and hence this condition guarantees that many intervals are occupied. In this case the law of large numbers applies and hence the energy becomes deterministic in the limit.

If $\lambda\nu = O(1)$, on the other hand, the value of $e_\omega(\gamma, \sigma, \nu)$ is random. This shows, in particular, that our condition on γ is optimal.

Also the condition $\sigma \gg \nu/(1 + \ln(1 + \nu^2/\gamma))$ can be expected to be optimal. It can be rephrased as $\bar{\ell}\sigma \gg 1$, where $\bar{\ell}$ is the (weighted) average length of occupied intervals

$$(21) \quad \bar{\ell} = \nu \int_0^\infty dP_\nu(\ell) \ell n(\ell)$$

with $n(\ell)$ the optimal repartition of the mass. A simple calculation shows that $\bar{\ell} \sim \nu^{-1}(1 + \ln(1 + \nu^2/\gamma))$.

5. CONCLUSIONS

- BEC in the ground state of the interacting gas in the GP regime can survive even in a strong random potential. The character of the wave function of the condensate, however, is strongly affected by the interaction.
- A random potential may lead to localization of the wave function of the condensate in subintervals. The interparticle interaction counteracts this effect, however, and can lead to complete delocalization (the condensate extends over the whole unit interval) if the interaction is strong enough.
- In terms of the interaction strength, γ , and density of scatterers, ν , the transition between localization and delocalization occurs in the model considered when $\gamma \sim \nu^2$. For $\gamma \lesssim \nu/(\ln \nu)^2$ a “third phase” occurs where the condensate is localized in a small number of subintervals.

REFERENCES

- [1] L.A. Pastur, A. Figotin, *Spectra of Random and Almost-Periodic Operators*, Springer-Verlag, 1992
- [2] O. Lenoble, L.A. Pastur, V.A. Zagrebnov, *Bose-Einstein condensation in random potentials*, C.R. Physique **5**, 129–142 (2004)
- [3] O.Lenoble, V.A.Zagrebnov, *Bose-Einstein Condensation in the Luttinger-Sy Model*, Markov Proc.Rel.Fields **13**, 441–468 (2007).
- [4] Th.Jaek, J.V.Pulé, V.A.Zagrebnov, *On the Nature of Bose-Einstein Condensation Enhanced by Localization*, J.Math.Phys. **51**, 103302-15 (2010).
- [5] H. Gimpferlein, S. Wessel, J. Schmiedmayer, L. Santos, *Ultracold Atoms in Optical lattices with Random One-Site Interactions*, Phys. Rev. Lett. **95**, 170401 (2005)
- [6] V.I. Yukalov, R. Graham, *Bose-Einstein condensed systems in random potentials*, Phys. Rev. A **75**, 023619 (2007)
- [7] P. Lugan, P. Bouyer, A. Aspect, M. Lewenstein, L. Sanches-Palencia, *Ultracold Bose Gases in 1D Disorder: From Lifshits Glass to Bose-Einstein Condensate*, Phys. Rev. Lett. **98**, 170403 (2007)
- [8] L. Fallani, C. Fort, M. Inguscio, *Bose-Einstein Condensates in Disordered Potentials*, Adv. At. Molec. Opt. Phys. **56**, 119–159 (2008)
- [9] L. Sanches-Palencia, D. Clément, P. Lugan, P. Bouyer and A. Aspect, *Disorder-induced trapping versus Anderson localization in Bose-Einstein condensates expanding in disordered potentials*, New J. Phys. **10**, 045019 (2008)
- [10] A.S. Pikovsky, D.L. Shepelyansky, *Destruction of Anderson Localization by a Weak Nonlinearity*, Phys. Rev. Lett. **100**, 094101 (2008)
- [11] P. Lugan, A. Aspect, L. Sanches-Palencia, D. Delande, B. Grémaud, C.A. Müller, C. Miniatura, *One-dimensional Anderson localization in certain correlated random potentials*, Phys. Rev. A **80**, 023605 (2009)
- [12] J. Radic, V. Bacic, D. Judic, M. Segev, H. Buljan, *Anderson localization of a Tonks-Girardeau gas in potentials with controlled disorder*, Phys. Rev. A **81**, 063639 (2010)
- [13] I.L. Aleiner, B.L. Altshuler, G.V. Shlapnykov, *A finite-temperature phase transition for disordered weakly interacting bosons in one dimension*, Nature Phys. **6**, 900–904 (2010)
- [14] L. Sanches-Palencia, M. Lewenstein, *Disordered quantum gases under control*, Nature Phys., **6**, 87–95 (2010)
- [15] M. Piraud, P. Lugan, B. Bouyer, A. Aspect, and L. Sanches-Palencia, *Localization of a matter wave packet in a disordered potential*, Phys. Rev. A **83** 031603(R) (2011)
- [16] W.B. Cardoso, A.T. Avelar, D. Bazeia, *Anderson localization of matter waves in chaotic potentials*, Nonlin. Analysis **13**, 755–763 (2012)
- [17] E.H. Lieb, W. Liniger, *Exact Analysis of an Interacting Bose Gas. I.*, Phys. Rev. **130**, 1605–1616 (1963)

- [18] E.H. Lieb, R. Seiringer, *Proof of Bose-Einstein Condensation for Dilute Trapped Gases*, Phys. Rev. Lett. **88**, 170409 (2002)
- [19] W. Kirsch, B. Simon, *Universal lower bounds on eigenvalue splittings for one dimensional Schrödinger operators*, Commun. Math. Phys. **97**, 453–460 (1985)

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY., 805 SHERBROOKE STREET WEST, MONTREAL, QC H3A 2K6, CANADA, E-MAIL: ROBERT.SEIRINGER@MCGILL.CA

FAKULTÄT FÜR PHYSIK, UNIVERSITÄT WIEN., BOLTZMANNGASSE 5, 1090 VIENNA, AUSTRIA, E-MAIL: JAKOB.YNGVASON@UNIVIE.AC.AT

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D'AIX-MARSEILLE (AMU) AND CENTRE DE PHYSIQUE THÉORIQUE - UMR 7332, LUMINY CASE 907, 13288 MARSEILLE, CEDEX09, FRANCE, E-MAIL: VALENTIN.ZAGREBNOV@UNIV-AMU.FR